

Incidence:
Measures the frequency with which an event occurs in a population over a time period

$$\text{Incidence} = \frac{\text{\# of new cases over time}}{\text{Population at risk over the same time period}}$$

$$\text{Prevalence} = \frac{\text{\# of cases}}{\text{Total population}}$$

Prevalence:
Measures what proportion of the population is affected

Measures of Disease Frequency

2 x 2 Tables

		Outcome		Total
Exposure	Yes	No		
	a	b	a + b	c + d
Yes	a	b	a + b	c + d
No	c	d	c + d	a + b + c + d
Total	a + c	b + d	a + b + c + d	a + b + c + d

a = number who are exposed and have the outcome
b = number who are not exposed and have the outcome

c = number who are not exposed and have the outcome
d = number who are not exposed and do not have the outcome

p = probability of having the outcome
q = probability of not having the outcome

$a + b + c + d = \text{total study population}$
 $a + b + c = \text{exposed population}$
 $a + c = \text{outcome population}$
 $b + d = \text{non-exposed population}$
 $c + d = \text{non-outcome population}$

$a + b = \text{total number who are exposed}$
 $c + d = \text{total number who are not exposed}$

$a + c = \text{total number who have the outcome}$
 $b + d = \text{total number who do not have the outcome}$

$a = \text{number who are exposed and have the outcome}$
 $b = \text{number who are not exposed and do not have the outcome}$

$c = \text{number who are not exposed and have the outcome}$
 $d = \text{number who are not exposed and do not have the outcome}$

Measures of Association

Assess the strength of association between an exposure and an outcome

$$\text{Relative Risk (RR)} = \frac{\frac{a}{c}}{\frac{a + b}{c + d}}$$

$$\text{Odds Ratio (OR)} = \frac{\frac{a * d}{c * b}}{\frac{a + c}{b + d}}$$

Interpreting RR and OR:

- = 1 indicates no association
- > 1 indicates a positive association
- < 1 indicates a negative association

Tests of Significance

Indicate the reliability of the association

95% Confidence Intervals (CI):


A range of possible values for the measure of association (RR/OR) that has a 95% chance of containing the true measure

- Range below 1.0 indicates less risk of outcome in exposed population
- Range above 1.0 indicates greater risk of outcome in exposed population

p-values:

Indicate how likely it would be for the observed measure (RR/OR) to occur by chance in absence of a true association

- Small p-values (< 0.05) indicate that the observed measure most likely was due to a true association between exposure and outcome



Source: Grimes DA, Schulz KF. Lancet 2002;359:58.



The basics of epidemiology for non-epidemiologists

Epidemiology Pocket Guide




The North Carolina Center for Public Health Preparedness is a program of the North Carolina Institute for Public Health in the University of North Carolina School of Public Health.

This publication was supported in part by grant/cooperative agreement number U90/CCU424265 from CDC. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC.

Basic epidemiology information... at your fingertips

Classification of Study Designs

Indicate the reliability of the association



Measures of Disease Frequency

2 x 2 Tables

Tests of Significance

Indicate the reliability of the association

95% Confidence Intervals (CI):


A range of possible values for the measure of association (RR/OR) that has a 95% chance of containing the true measure

- Range below 1.0 indicates less risk of outcome in exposed population
- Range above 1.0 indicates greater risk of outcome in exposed population

p-values:


Indicate how likely it would be for the observed measure (RR/OR) to occur by chance in absence of a true association

- Small p-values (< 0.05) indicate that the observed measure most likely was due to a true association between exposure and outcome



Classification of Study Designs

Indicate the reliability of the association



Measures of Disease Frequency

Indicate the reliability of the association

95% Confidence Intervals (CI):

A range of possible values for the measure of association (RR/OR) that has a 95% chance of containing the true measure

- Range below 1.0 indicates less risk of outcome in exposed population
- Range above 1.0 indicates greater risk of outcome in exposed population

p-values:

Indicate how likely it would be for the observed measure (RR/OR) to occur by chance in absence of a true association

- Small p-values (< 0.05) indicate that the observed measure most likely was due to a true association between exposure and outcome